



MYCO ACTIVE

Soil bio-stimulant to activate beneficial microbes



pH 3.5

SG 1.23

Product	N	P	K	S	Ca	Mg	Fe	Si	C	Zn	Mn	Cu	Mo	B	Co	HUMIC ACID	FULVIC ACID
	%	%	%	%	%	%	%	%	%	ppm	ppm	ppm	ppm	ppm	ppm	%	%
MYCO ACTIVE	3.2	1.2	2.8	3.3		0.18			0.4	4,537	7,323	131	8	135	4		0.14

Typical uses and benefits include:

- Boosts plant nutrient and moisture uptake
- Improves root development
- Increases plant vigour
- Crop establishment

Ecogrowth's Myco Active enhances nutrient and moisture uptake by emulating root exudates and stimulating beneficial microbes.

Myco Active contains a complex range of compounds that perform or mimic the function of natural hormone-like biochemicals, diffusing through the soil from the root system. These compounds stimulate responses from the soil indigenous microbial population to balance/enhance activity.

COMPLEMENTARY PRODUCTS



ECO-HUMATE

Concentrated liquid humus soil conditioner.



ECO-VITAL®

A premium natural kelp and fish liquid fertiliser.



potassium SILICATE

Premium silica & potassium liquid fertiliser.

fertiliser for life

AVAILABLE IN 5L, 20L, 200L, IBC & BULK

For alternative applications or to arrange a soil/plant test please contact your Ecogrowth® nutritional expert.



MYCO ACTIVE



Soil bio-stimulant to activate beneficial microbes

APPLICATION RATES

Crop	Rate	Notes
TREE CROPS (Pome, Stone, Nuts, Mangoes, Avocados)	Establishment: 20ml/plant Start Of Season: 3-6L/ha soil/fertigation Support: 1-2L/ha soil/fertigation 1:200 min. dilution	Apply to soil at budburst, then monthly for support.
VINES WINE & TABLE	Establishment: 20ml/plant Start Of Season: 2-4L/ha soil/fertigation Support: 1-2L/ha soil/fertigation 1:200 min. dilution	Apply to soil at budburst, then monthly for support.
VEGETABLES	Start Of Season: 4-8L/ha soil/fertigation Support: 2-4L/ha soil/fertigation 1:200 min. dilution	Apply throughout season.
TURF	5-15L/ha 50-150ml/100m ² 1:80 min. dilution	Water in for best results.
BROADACRE & PASTURE	2+L/ha foliar 5+L/ha furrow 1:40 min. dilution	Apply as required. In furrow treatment should be banded away from seed at lower water rates.
NURSERY	3ml/m ² Fertigated	Apply at establishment, re-potting and after transplant.